

Oral Medicine and Surgical Sciences Issue (Oral Medicine, Oral and Maxillofacial Surgery, Oral Pathology, Oral Biology)

4-1-2020

## Evaluation of Topical Application of Nigella Sativa (Black Seeds) on Delayed Dental Implant

Zahraa Ahmed

*Demonstrator at Oral medicine , Periodontology , Oral Diagnosis and Radiology Department, Faculty of Dental Medicine for Girls, Al-Azhar University, Cairo, Egypt, hos142000@yahoo.com*

Mai Attia

*Assistant Professor of Oral Medicine, Periodontology, Oral Diagnosis and Radiology Department, Faculty of Dental Medicine for Girls , Al-Azhar University, Cairo, Egypt, mai\_shafik@yahoo.com*

Aya Mahmoud

*Lecturer of Pharmaceutics, Faculty of pharmacy for Girls, Al- Azhar University, Cairo, Egypt, ayadawaba@azhar.edu.eg*

Eatemad Shoreibah

*Professor of Oral Medicine, Periodontology, Oral Diagnosis and Radiology Department, Faculty of Dental Medicine for Girls, Al-Azhar University, Cairo, Egypt, eshoreibah@yahoo.com*

Follow this and additional works at: <https://azjd.researchcommons.org/journal>



Part of the [Other Dentistry Commons](#)

---

### How to Cite This Article

Ahmed, Zahraa; Attia, Mai; Mahmoud, Aya; and Shoreibah, Eatemad (2020) "Evaluation of Topical Application of Nigella Sativa (Black Seeds) on Delayed Dental Implant," *Al-Azhar Journal of Dentistry*: Vol. 7: Iss. 2, Article 5.

DOI: <https://doi.org/10.21608/adjg.2020.7568.1082>

This Original Study is brought to you for free and open access by Al-Azhar Journal of Dentistry. It has been accepted for inclusion in Al-Azhar Journal of Dentistry by an authorized editor of Al-Azhar Journal of Dentistry. For more information, please contact [yasmeenmahdy@yahoo.com](mailto:yasmeenmahdy@yahoo.com).



## Evaluation of Topical Application of Nigella Sativa (Black Seeds) on Delayed Dental Implant

Zahraa M. Ahmed <sup>1\*</sup>, Mai S. Attia <sup>2</sup>, Aya M. Mahmoud <sup>3</sup> and Eatemad A. Shoreibah <sup>4</sup>

Codex : 32/2004

azhardentj@azhar.edu.eg

<http://adjg.journals.ekb.eg>

DOI: 10.21608/adjg.2020.7568.1082

Oral Medicine & Surgical Sciences  
(Oral Medicine, Oral & Maxillofacial  
Surgery, Oral Pathology, Oral Biology)

### ABSTRACT

**Purpose:** The present study was conducted to evaluate the effect of topically applied Nigella Sativa (Black Seeds) on osseointegration of delayed dental implant in an effort to improve outcomes of dental implants. **Subjects and Methods:** The current study was performed on twelve (12) healthy patients that were divided randomly into two groups: Group 1 (Test group) included six (6) patients with delayed implant insertion with the topical use of nigella sativa; Group 2 (Control group) included six (6) patients with delayed implant insertion without the use of nigella sativa. **Results:** A non-statistically significant difference was recorded in clinical parameters after three and six months follow up in both groups. Test group showed a highly significant increase in its bone density after six months in comparison with control group. **Conclusion:** The topical use of nigella sativa in delayed dental implant showed an improvement in peri-implant tissues.

### INTRODUCTION

Dental implant is a valuable management approach in complete or partially edentulous patients, playing an important role in orofacial anatomic contour reconstruction, health, function, and esthetic <sup>(1,2)</sup>.

### KEYWORDS

*Dental implants, Nigella Sativa,  
Modified gingival index*

A paper extracted from Master Thesis titled "Evaluation of Topical Application of Nigella Sativa (Black Seeds) on Delayed Dental Implant"

1. Demonstrator at Oral medicine , Periodontology , Oral Diagnosis and Radiology Department, Faculty of Dental Medicine for Girls, Al-Azhar University, Cairo, Egypt
2. Assistant Professor of Oral Medicine, Periodontology, Oral Diagnosis and Radiology Department, Faculty of Dental Medicine for Girls , Al-Azhar University, Cairo, Egypt
3. Lecturer of Pharmaceutics, Faculty of pharmacy for Girls, Al- Azhar University, Cairo, Egypt
4. Professor of Oral Medicine, Periodontology, Oral Diagnosis and Radiology Department, Faculty of Dental Medicine for Girls, Al-Azhar University, Cairo, Egypt

\* Corresponding author email: hos142000@yahoo.com

About 450,000 osseointegrated dental implants are being placed annually, with high success rate reach 95% specially in case of solitary tooth replacement with an implant supported crown <sup>(3)</sup>.

The original purpose of the dental implant treatment and to guarantee implant success is to produce and maintain a firm connection between bone and dental implant. This idea has been explained as composed of highly distinguishable tissue making” a direct structural and functional connection between prepared, existing bone and the surface of a load carrying implant”<sup>(4)</sup>.

Bone quality is one of the main factors that have an effect on osseointegration of dental implant, reduced bone quality and/or quantity, believed as one of the most significant causes of implant failure as it leading to unfortunate anchorage and stability. Bone mass relays on the level of bone density there. Bone quality can progress in the region of a functional osseointegrated dental implant as a result of the positive bone stimulation, the extra bone that is present by the side of an implant place, the enhanced chance for implant success <sup>(5)</sup>.

Implant surface alteration has been considered and used to enhance natural surface properties helping osseointegration <sup>(6)</sup>. The main issue in implant osseointegration is surface irregularity, that demonstrated to raise osteoblast action in contrast to a smooth surface <sup>(7)</sup>. Surface roughness of the implant has been improved via different means as machining, plasma spray coating, grit blasting, acid etching, sandblasted and acid etching (SLA), anodizing, and biomimetic coating <sup>(8,9)</sup>.

With the purpose of encourage wound healing and attain an optimized osseointegration, systemic and local administration of drugs, plus growth and differentiation factors and / or implant drug based surface alterations have been engaged <sup>(10,11)</sup>. In the older years, the text concerning the local interferences become highly developed <sup>(11)</sup>.

World is rising attention to treat different diseases by using medicinal herbs or plants due to their therapeutic potential and less side effects <sup>(12)</sup>.

One of the most classified as evidence based herbal medicine is *Nigella Sativa* (NS) <sup>(13,14)</sup>. *Nigella Sativa* is a flower plant within the family Ranunculaceae resident to the south and southwest Asia, and is developed in numerous countries within the Mediterranean area as South Europe, Syria, Turkey and Saudi Arabia. *Nigella Sativa* also recognized as black seed, black cumin, the seed of bleesing or Habbatul Barakah <sup>(15,16)</sup>.

The black seed has a religious and spiritual energy for the Muslims, as its therapeutic use was started after the arrival of Islam since **prophet Mohammed (peace and pray upon him)** talk about its therapeutic value and likely of cure, when said that” there is cure for every disease in black seed except death “ <sup>(17)</sup>. This give explanation why NS is broadly used by many Muslims all over the world <sup>(18)</sup>.

The pharmacological action of the NS expose a broad spectrum of activity including antidiabetic<sup>(19,20)</sup>, antihypertensive, anticancer, immunomodulator<sup>(20)</sup>, antimicrobial, anti-inflammatory, analgesic, antihistaminic, bronchodialator, liver protection, kidney protection, spasmolytic and antioxidant qualities <sup>(21)</sup>.

On the topic of implant dentistry, in an invivo study, the black seed oil extract was utilized to inspect its outcome on the bone implant interface’s (mechanical and histological properties). Inside this experimental study, 12 New Zealand rabbits take delivery of 4 screw-shaped implants in the tibiae bone, the implant were separated into 2 groups; coated and uncoated with black seed oil extract, over a period of time. the end results demonstrated that, mechanically the coated implants had considerably superior torque resistance than that of uncoated implants, and histologically revealed early osseointegration with an osteophilic surface, besides, no undesirable tissue reaction <sup>(22)</sup>.

CBCT applied in implant dentistry to estimate the bone quality and quantity and to identify any bone diseases at implant site <sup>(23,24)</sup> and it was used in this study to estimate bone density changes before and after implant insertion.

## SUBJECTS AND METHODS

### Study design

The current study was a randomized clinical trial on 12 patients selected from the out-patient Clinic of Oral Medicine & Periodontology Department, Faculty of Dental Medicine for Girls, Al-Azhar University.

### Sample Size:

Sample size calculations achieved using <http://biomath.info/power> according to the study (Clinical and Radiographic Evaluation of Immediate Loaded Dental Implants with Local Application of Melatonin: A Preliminary Randomized Controlled Clinical Trial)<sup>(25)</sup>. A total sample size of 12 patients (6 patients in each group) was sufficient to identify the difference. Total numbers of patients were divided randomly in to two groups;

Group 1 (Test group): 6 patients with delayed implant placement with the topical use of nigella sativa.

Group 2 (Control group): 6 patients with delayed implant placement without using nigella sativa

The patients were selected according to selected criteria (completely healed surgical site, presences of proper inter arch space for the placement of the implant prosthetic part, bone density ranging from D2 to D3, non-smoker and non-pregnant women). The selected patients signed an informed consent explaining all the procedures to the patients including all benefits and side effects in simple and easy way.

### Surgical protocol:

After local anesthesia administration, crestal incision at the edentulous site with sulcular

incisions around mesial and distal natural tooth were performed. Full thickness flap was elevated using mucoperiosteal elevator. Osteotomy site preparation through sequential drilling followed the protocol described by the implant company surgical kit (Dentium super line implant system (Emergo Europe, Seoul, Korea)) until reaching the desired diameter of the implant under copious saline irrigation, then the implant screwed in clockwise manner until complete seating of the implant to its final insertion depth. Flap was returned to its position covering the implant and sutured with 4-0 polypropylene (vicryl, absorbable) suture material.

Regarding test group (group1), the same steps were followed but, after preparation of osteotomy site and before implant placement, Nigella Sativa gel was applied inside the prepared site until the gel fill the whole osteotomy site. Augmentin 1g tablet antibiotic was prescribed twice per day for at least 5 days after surgery, Ibuprofen 400 mg tablet was given twice daily for 3 or 5 days as analgesic and anti-inflammatory, Chlorhexidine 0.12% mouthwash was prescribed twice daily for one week after surgery.

Sutures were removed 10 days postoperatively. Two months after surgery, the cover screw of the implant was removed and gingival former was placed.

**Nigella Sativa gel preparation:** the gel prepared under complete aseptic condition according to this paper <sup>(26)</sup>.

### Clinical evaluation:

Modified Gingival index (MGI) was recorded for each patient at 3-months and 6- months after implant placement.

Probing depth (PD): It was obtained by measuring the probing pocket per site around the implant. Probing depth is measured from the gingival margin to the base of the sulcus. Probing depth was recorded at 3 and 6 months for each patient around the healing abutment.

### Radiographic Assessment:

Cone beam computed tomography (CBCT) was used twice before fixture placement and at the end of the study (after 6 months) to record bone density.

Bone density was taken for the sagittal and coronal view. For each view, two lines were drawn along the total length of the implant fixture (mesial and distal aspect of the sagittal view, buccal and palatal of coronal view) and then each line splitted into three parts, corresponding the coronal, middle and apical thirds. Measurements were obtained from around approximately 1 mm in a parallel manner away from the implant fixture to avoid titanium artifact at the bone implant interface. Then these six readings were divided by six to get the mean value of bone density around each implant in each view. The readings were recorded in Hounsfield Units (HU) <sup>(27)</sup>.

### STATISTICAL ANALYSIS

Data were collected, revised, coded and entered to the Statistical Package for Social Science

(IBM SPSS) version 23. The quantitative data were presented as mean, standard deviations and ranges while qualitative variables were presented as number and percentages.

### RESULTS

#### Clinical evaluation results:

Modified Gingival Index: at 3 months, test group showed a lower MGI mean ( $0.04 \pm 0.10$ ) than control group ( $0.13 \pm 0.21$ ), however this difference was statistically insignificant ( $P=0.401$ ). At 6 months, test group showed a lower MGI mean ( $0.08 \pm 0.20$ ) than control group ( $0.21 \pm 0.29$ ), however this difference was statistically insignificant ( $P=0.411$ ) (table 1).

Probing Depth: at 3 months, test group showed a lower PD mean ( $0.96 \pm 0.68$ ) than that of control group ( $1.83 \pm 0.70$ ), however this difference was statistically insignificant ( $P=0.053$ ). At 6 months, test group showed a lower PD mean ( $0.63 \pm 0.16$ ) than control group ( $1.44 \pm 1.05$ ), however this difference was statistically insignificant ( $P=0.090$ ) (table 1).

**Table (1):** Results of MGI and PD changes of both groups through the study period:

Mean of M.G.I after implant insertion		Control group	Test group	Test value	P-value	Sig.
		No. = 6	No. = 6			
At 3 months	Mean $\pm$ SD	$0.13 \pm 0.21$	$0.04 \pm 0.10$	0.877	0.401	NS
	Range	0 – 0.5	0 – 0.25			
At 6 months	Mean $\pm$ SD	$0.21 \pm 0.29$	$0.08 \pm 0.20$	0.859	0.411	NS
	Range	0 – 0.75	0 – 0.5			
Paired t T-test	t	1.000	1.000			
	p-value	0.363 (NS)	0.363 (NS)			
Mean of P.D after implant insertion		Control group	Test group	Test value	P-value	Sig.
		No. = 6	No. = 6			
At 3 months	Mean $\pm$ SD	$1.83 \pm 0.70$	$0.96 \pm 0.68$	2.197	0.053	NS
	Range	0.75 – 2.5	0.5 – 2.25			
At 6 months	Mean $\pm$ SD	$1.44 \pm 1.05$	$0.63 \pm 0.16$	1.878	0.090	NS
	Range	0.5 – 3	0.5 – 0.88			
Paired t-test	t	1.290	1.384			
	p-value	0.254 (NS)	0.225 (NS)			

$P > 0.05$ : Non significant ,  $P < 0.05$ : Significant,  $P < 0.01$ : Highly significant

**Radiographic assessment results:**

Bone density (coronal view): Within test group there was increase in the bone density reading before implant insertion and after 6 months ( $713.38 \pm 124.37$  and  $969.87 \pm 182.13$ ) respectively; with high significant difference ( $P=0.002$ ) (table 2) .

Bone density (sagittal view): At 6 months after implant insertion; control group and test group

readings were  $531.18 \pm 76.13$  and  $795.65 \pm 127.35$  respectively. This demonstrated high statistically significant difference ( $P=0.001$ ). Within test group there was increase in the bone density reading before implant insertion and after 6 months ( $627.74 \pm 127.53$  and  $795.65 \pm 127.35$ ) respectively; with high statistically significant difference ( $P=0.005$ ) (table 2) .

**Table (2):** Results of bone density (coronal and sagittal view) changes of both groups through the study period .

Mean of bone density (coronal view)		Control group	Test group	Test value	P-value	Sig.
		No. = 6	No. = 6			
Before implant insertion	Mean $\pm$ SD	731.88 $\pm$ 130.89	713.38 $\pm$ 124.37	0.251	0.807	NS
	Range	488.95 – 860.95	506.1 – 820.8			
At 6 months after implant insertion	Mean $\pm$ SD	799.60 $\pm$ 181.29	969.87 $\pm$ 182.13	-1.623-	0.136	NS
	Range	641.8 – 1073.95	629.65 – 1181			
Paired t-test	t	0.887	5.886			
	p-value	0.416 (NS)	0.002 (HS)			
Mean of bone density (sagittal view)		Control group	Test group	Test value	P-value	Sig.
		No. = 6	No. = 6			
Before implant insertion	Mean $\pm$ SD	659.44 $\pm$ 87.13	627.74 $\pm$ 127.53	0.503	0.626	NS
	Range	557.65– 806.1	441.3 – 752.3			
At 6 months after implant insertion	Mean $\pm$ SD	531.18 $\pm$ 76.13	795.65 $\pm$ 127.35	-4.366-	0.001	HS
	Range	443.1– 664.95	596.95 – 924.95			
Paired t-test	t	2.275	4.882			
	p-value	0.072(NS)	0.005 (HS)			

$P > 0.05$ : Non significant ,  $P < 0.05$ : Significant,  $P < 0.01$ : Highly significant

**DISCUSSION**

Criteria to assess the success and failure of dental implants which involves clinical parameters (Modified Gingival index and probing depth) and radiographic measurement was carried out <sup>(28)</sup>. As regards Modified gingival index <sup>(29)</sup>, it measured to express disease activity <sup>(30)</sup>. On the other hand, probing depth index is a critical index for identification of peri-implant disease <sup>(31)</sup> .

In terms of clinical parameters (MGI and PD) in this study, no significant difference was recorded between both groups at 3 and 6 months postoperative following implant insertion, this could be attributed to adequate patient motivation and proper oral hygiene <sup>(32)</sup>.

Regarding bone density results; Before implant insertion no statistically significant was found between both groups ( $P=0.626$ ), while, there was



highly statistically significant difference ( $P=0.001$ ) 6 months after implant insertion.

The consequences of the improved bone around dental implant and enhance bone density in test group, this could be qualified to biological action of NS. Where, NS works as bioactive and bioinductive substance that stimulate and increase speed of bone formation this could be attributed to the existence of its components as Protein and amino acids. NS seed had 22.6% - 26.7% protein and amino acids mainly Glutamic acid, Arginine and Aspartic acid<sup>(33)</sup>. Moreover, rapid bone formation caused by black seed oil extract could be attributed to the presence of calcium and phosphorous in its chemical constituents which are essential for bone formation<sup>(34)</sup>.

## CONCLUSION

The use of *Nigella Sativa* gel in delayed dental implants may be able to improve bone quality around dental implants as it increase bone density, thus improving implant osseointegration which leading to good prognosis for a long-term survival.

## REFERENCES

- Beikler T, Flemmig TF. Implants in the medically compromised patient. *Crit Rev Oral Biol Med*. 2003;14:305-16.
- Chanavaz M. Patient screening and medical evaluation for implant and preprosthetic surgery. *J Oral Implantol*. 1998;24:222-9.
- Sullivan RM. Implant dentistry and the concept of osseointegration: a historical perspective. *J Calif Dent Assoc*. 2001;29:737-45.
- Branemark PI. Osseointegration and its experimental studies. *J Prosthet Dent*. 1983;50:399-410.
- McNutt MD, Chou CH. Current trends in immediate osseous dental implant case selection criteria. *J Dent Educ*. 2003;67:850-9.
- Smeets R, Stadlinger B, Schwarz F, Beck-Broichsitter B, Jung O, Precht C, modifications on osseointegration. *Biomed Res Int* 2016;6285620.
- Von Wilmsowky C, Moest T, Nkenke E, Stelzle F, Schlegel KA. Implants in bone: part I. A current overview about tissue response, surface modifications and future perspectives. *Oral Maxillofac Surg*. 2014; 18:243-57.
- Shemtov-Yona K, Rittel D. An overview of the mechanical integrity of dental implants. *Biomed Res Int*. 2015:547384.
- Jemat A, Ghazali MU, Razali M, Otsuka Y. Surface modification and their effect on titanium dental implants. *Biomed Res Int* 2015:791725.
- Dimitriou R, Jounes E, McGonagle D, Giannoudis PV. Bone regeneration: current concepts and future directions. *BMC Med*. 2011;9:66.
- Apostu D, Lucaciu GD, Crisan B, Crisan L, Baciut M, Onisor F, et al. Systemic drugs that influence titanium implant osseointegration. *Drug Metab Rev*. 2017; 49:92-104
- Abd-Awn B, Al-Dhaher Z, Al-Dafaai R. The effect of black seed oil extracts on mutans streptococci in comparison to chlorhexidine gluconate (in vitro). *J f Baghdad Coll Dent* 2012; 24:126-131.
- Ahmad I, Tripathi J, Sharma M, Karchulli MS, Umer L. *Nigella sativa* –a medicinal herb with immense therapeutic potential (a systematic review). *Int J Biol Pharma Res*. 2014; 5: 755-62.
- Ahmad A, Husian A, Mujeeb M, Khan S, Najmi A, Siddique N, et al. A review on therapeutic potential of *Nigella sativa*; a miracle herb. *Asian Pac J Trop Biomed* 2013; 3: 337-52.
- Tariq M. *Nigella sativa* feeds: folklore treatment in modern day medicine. *Saudi J Gastroenterol* 2008; 14: 105-6.
- Shrivastava R, Agrawal R, Parveen Z. A review on therapeutic applications of *Nigella Sativa*. *Journal of Chemistry and Chemical Sciences* 2011; 1: 241-8.
- AlAttas SA, Zahran FM, Turkistany SA. A review on *Nigella sativa* and its active constituent thymoquinone in oral health. *Saudi Med J*. 2016; 37:235-44.
- Rhamani AH, Alzhairy MA, Khan MA, Aly SM. *Nigella Sativa* and its active constituents thymoquinone shows pivotal role in the diseases prevention and treatment. *Asian J Pharm and Clin Res*. 2015; 8: 48-53.
- Hmza AJ, Osman MT, Anan A, Omar E. Immunomodulatory effect of *Nigella sativa* oil in the disease process of type 1 diabetic rats. *Res J Pharm Bio Chem Sci* 2013; 4: 980-8.
- Hmza AJ, Omar E, Anan A, Osman MT. *Nigella sativa* oil has significant repairing ability of damaged pancreatic tissue occurs in induced type 1 diabetes mellitus. *Glob J Pharmacol* 2013; 7: 14-9.

21. Ahmad A, Husian A, Mujeeb M, Khan S, Najmi A, Siddique N, et al. A review on therapeutic potential of Nigella sativa; a miracle herb. *Asian Pac J Trop Biomed* 2013; 3: 337-52.
22. Alnajjar SSA, Mohammed SA. Mechanical and histological significance of Nigella Sativa oil extract on bone-implant interface. *J. Baghdad Coll. Dent.* 2009; 21: 39-43.
23. Jacobs R, Mraiwa N, vanSteenberghe D, Gijbels F, Quirynen M. Appearance, location, course, and morphology of the mandibular incisive canal: an assessment on spiral CT scan. *Dento maxillofac Radiol.* 2003;31:322-7.
24. Guler AU, Sumer M, Sumer P, Bicer I. The evaluation of vertical heights of maxillary and mandibular bones and the location of anatomic landmarks in panoramic radiographs of edentulous patients for implant dentistry. *J Oral Rehabil.* 2005;32:741-6.
25. El-Gammal MY, Salem AS, Anees MM, Tawfik MA. Clinical and Radiographic Evaluation of Immediate Loaded Dental Implants with Local Application of Melatonin: A Preliminary Randomized Controlled Clinical Trial, *J. oral implantol.* 2016; 42:119-125.
26. Auda SH, Abd El-Rasoul S, Ahmed MM, Osman SK, Mahmoud El-Badry M. In-vitro release and in-vivo performance of tolmetin from different topical gel formulations. *J Pharm Investig* 2015; 45:311-7.
27. Elkhidir Y, Wei S, Suyang L, Xie M, Yang C. Feasibility of CBCT in Evaluating Bone Density of Dental Implant Placement Sites. *Research & Reviews: J. Dent. Sci.* 2017;5:1.
28. Salvi GE, Lang NP. Diagnostic parameters for monitoring peri-implant conditions. *Int J Oral Maxillofac Implants* 2004;19:116-27.
29. Mombelli A, Van Oosten MA, Schurch E and Land NP. The microbiota associated with successful or failing osseointegrated titanium implants. *Oral Microbial Immunol* 1987; 2:145-51.
30. Sebastian GC. Current status of indices of gingivitis. *J. Clin. Periodontol.* 1986; 13: 375-8.
31. Lang NP, Bragger U, Walther D, Beamer B and Kornman K. Ligature-induced peri-implant infection in cynomolgus monkeys. I. Clinical and radiographic findings. *Clin. Oral Implants Res.* 1993; 4: 2-11.
32. Abraham HM, Philip JM, Kruppa J, Jain AR and Krishnan CJV. Use of Chlorhexidine in Implant Dentistry. *Biomed Pharmacol J* 2015;8:1-3.
33. Al-Jassir SM. Chemical composition and microflora of black cumin (*Nigella Sativa* L.) seeds growing in Saudi Arabia. *Food Chem.* 1992; 45: 239-42.
34. Al-Hijazi AY, Mohammed HS. Evaluation of the effect of Nigella Sativa oil and powder on socket healing process. *J Nat Sci Res* 2013; 3: 135-41.